

A Case Report of Giant Tonsillolith

Smitha Soubhagya Gangaraj, Nirmala Maruthi

ABSTRACT

Tonsilloliths, also known as tonsil stones or tonsillar calculi, are clusters of calcified material that form in the crypts of palatine tonsils. They are usually of small size. Large or giant tonsilloliths are rare. We report a case of giant tonsillolith in the right palatine tonsil and literature is reviewed.

Keywords: Tonsillolith, Tonsils, Calculus.

Source of support: Nil

Conflict of interest: None declared

INTRODUCTION

Tonsilloliths are calcified concretions that develop in tonsillar crypts, within the substance of the tonsil or around it. They occur in up to 10% of the population, frequently due to episodes of tonsillitis. These calculi are composed of calcium salts either alone or in combination with other mineral salts. Small concretions are relatively common but large or giant tonsilloliths are rare¹ and very few cases are reported in the literature.² We report a case of giant tonsillolith in the right palatine tonsil.

CASE REPORT

A 56-year-old female patient presented with foreign body sensation in the throat since 3 months. It was associated with halitosis and occasional right ear ache. She had recurrent episodes of throat pain in childhood.

Oral examination revealed a large, brownish-black mass protruding from the right tonsillar tissue mimicking a foreign



Fig. 2: Brownish-black tonsillolith measuring 2 × 1.5 cm

body (Fig. 1). On palpation, the mass was stony hard in consistency with irregular rough surface.

The tonsillolith measuring about 2 × 1.5 cm was removed under topical anesthesia (Fig. 2) and oral mouth gargle was prescribed. The patient is now asymptomatic without any recurrence.

DISCUSSION

The earliest known description of oropharynx concretions is recorded in 1560.¹ Tonsilloliths are calcified structures of bacteria and organic debris that commonly develop in the tonsillar crypts, within the substance of the tonsil or around it. They occur between 20 and 77 years of age and are rare in pediatric age group.³ In a review by Mesoellella et al,² they were found to be located in the tonsillar tissue in 69.7%, in the tonsillar fossa in 21.2 and 9% were palatine in location. Only one case of calculi in the lingual tonsil has been reported in the literature.⁴

The exact pathogenesis of these calculi is unknown although there are many hypothesis on the formation of these. It has been stated that they originate as a result of repeated tonsillitis which lead to fibrosis of the ducts of crypts and retention of epithelial debris thereof. This epithelial debris forms the ideal media of growth of bacterial, actinomyces and fungi, such as *Leptothrix buccalis*. Finally, dystrophic calcification occurs as a result of deposition of inorganic salts from the saliva secreted by major and minor salivary glands.⁵ Other authors have proposed alternative



Fig. 1: Tonsillolith embedded in the right tonsil

mechanisms when the calculi are located in the peritonsillar areas, such as existence of ectopic tonsillar tissue, the formation of calculi secondary to salivary stasis within the minor salivary gland ducts in these locations or the calcification of abscessified accumulation.⁶

Clinical signs and symptoms are usually absent in small tonsilloliths. Larger ones may present with foreign body sensation in the throat, recurrent halitosis, odynophagia and referred otalgia. On throat examination, tonsillolith may be seen embedded in the tonsillar crypt. They may be single or multiple and of variable shapes like round, oval, cylindrical, pyramidal or plurilocular. The color also varies from grayish-yellow to dark gray, black or red brown.² It may present as tonsil or tonsillar fossa swelling when it is embedded inside the tonsillar tissue or fossa and can be palpated as a hard mass.⁷ The diagnosis can be easily made by clinical examination. Doubtful cases can be confirmed with imaging diagnostic techniques which will show the radiopaque shadow. Differential diagnosis of tonsilloliths includes foreign body, calcified granuloma, malignancy, an enlarged styloid process or rarely isolated bone which is usually derived from embryonic rests originating from the branchial arches.⁸

Treatment usually involves removal of tonsillolith under topical or local anesthesia. Vigorous mouth gargling daily helps in keeping the tonsil crypts clear of debris. In case of a large tonsillolith impacted within the tonsil or in chronic tonsillitis, tonsillectomy is indicated.⁹ In recurrent tonsilloliths, laser cryptolysis can be done to decrease the surface area of the tonsils via laser surfacing. A scanned carbon dioxide laser selectively vaporizes and smoothens

the surface of the tonsils and thus flattens the edges of crypts preventing trapped material from forming calculus.

REFERENCES

1. Pruet CW, Duplan DA. Tonsil concretions and tonsilloliths. *Otolaryngol Clin North Am* 1987;20:305-09.
2. Mosella M, Cimmino M, Martino DM, et al. Tonsillolith: A case report and review of the literature. *Acta Otorhinolaryngol Ital* 2004;24:302-07.
3. Cooper MM, Steinberg JJ, Lastra M, Antopol S. Tonsillar calculi: Report of a case and review of the literature. *Oral Surg Oral Med Oral Pathol* 1983;55:239-43.
4. Bando H, Uno T, Nin F, Tie K, Shinomiye T, Hisa Y. A case of calculus of the lingual tonsil. *Practica Otorhinolaryngologica* 2003;97(7):239-43.
5. Paparella MM, Shumrick DA. *Otolaryngology (Vol 3, Head & neck (3rd ed))*. WB Saunders: Philadelphia 1991:2141.
6. Revel MP, Laccocureye O, Hartl D, Bely N, Nando P, Brasnu D. Giant tonsillolith. *Ann Otol Laryngol* 1998;107:262-63.
7. Sezer B, Tugsel Z, Bilgen C. An unusual tonsillolith. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;95:471-73.
8. Aspestrand F, Kolbenstvedt A. Calcifications of the palatine tonsillary region: CT demonstration. *Radiology* 1987;165:479-80.
9. Silvestre FJ, Pla-Mocholi A, Estelles FE, Martinez MV. Giant tonsillolith: Report of a case. *Med Oral Pathol Oral Cir Bucal* 2005;10(3):239-42.

ABOUT THE AUTHORS

Smitha Soubhagya Gangaraj (Corresponding Author)

Assistant Professor, Department of ENT, Kempegowda Institute of Medical Sciences, Bengaluru, Karnataka, India, e-mail: smitha_praveen@hotmail.com

Nirmala Maruthi

Senior Resident, Department of ENT, Kempegowda Institute of Medical Sciences, Bengaluru, Karnataka, India